



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Merganser and there is a final chapter on conclusions in the matter of the extension of the breeding range in ducks.

Four of the chapters have already appeared in 'British Birds' but the others are here published for the first time. The work forms a valuable contribution to our knowledge of the Anatidae.—W. S.

Chapman on South American Birds.—In a paper on the genus *Pygochelidon*,¹ Dr. Chapman recognizes *P. cyanoleuca* of rather wide range in the mountains, *P. patagonica patagonica* of the South Temperate zone, *P. p. peruviana* (p. 7) a new race from Peru, and *P. flavipes* (p. 8) a very distinct species represented by one specimen from Prov. Juvien, Peru. Interesting comments on the probable origin and distribution of the forms follow. In another contribution² Dr. Chapman describes eight new races and species from Colombia, Ecuador and Argentina including *Jacana scapularis* (p. 3) from western Ecuador, the first record of the genus from west of the Andes.—W. S.

Wetmore and Peters on New American Birds.³—*Teledromas* (p. 41) is proposed for *Rhinocrypta fusca* Scl. and Salv. *Dendrocygna bicolor helva* (p. 42), San Diego Co., Calif., is described as new, the North American bird proving to be separable from true *bicolor* of Paraguay. *Colaptes pitius cachimans*, (p. 43) Argentina, is proposed as new and the genera *Pituipicus* and *Soroplex* are merged in *Colaptes*. *Brachyspiza capensis choraules* (p. 44), Rio Negro and *Saltator aurantirostris nasica* (p. 45) Mendoza, Argentina, are described, and Dr. Wetmore independently describes *B. c. mellea* (p. 39) from Paraguay⁴ and three new forms of Tinamou;⁵ *Rhynchotus arcanus* (p. 434). Parana, Argentina; *Nothura maculosa savannarum* (p. 435) Rocha, Uruguay; and *Calopezus elegans albidus* (p. 437), San Juan, Argentina.—W. S.

Bangs on Philippine Birds.⁶—This paper consists of comments on eight species and descriptions of five new forms all from extensive collections presented to the Museum of Comparative Zoology by Governor General Forbes. The new races are *Lalage niger mitifica* (p. 80) Lubang; *Aegithina*

¹ The Distribution of the Swallows of the Genus *Pygochelidon*. By Frank M. Chapman. Amer. Mus. Nov., No. 30, February 28, 1922, pp. 1-15.

² Descriptions of Apparently New Birds from Colombia, Ecuador, and Argentina. By Frank M. Chapman. March 2, 1922.

³ A New Genus and Four New Subspecies of American Birds. By Alexander Wetmore and James L. Peters. Proc. Biol. Soc., Wash., 35. pp. 41-46. March 20, 1922.

⁴ Description of a *Brachyspiza* from the Chaco of Argentina and Paraguay. *Ibid.* pp. 39-40, March 20, 1922.

⁵ Three new Birds of the Family Tinamidae. By Alexander Wetmore. Jour. Wash. Acad. Sci., 11, No. 18, Nov. 4, 1921.

⁶ Notes on Philippine Birds Collected by Governor W. Cameron Forbes. Bull. Mus. Comp. Zool. LXV., No. 4. April, 1922, pp. 77-84.